

PLEASE BRING THIS PACKET WITH YOU ON THE DAY OF SURGERY

Pre and Post Operation Instructions for Cosmetic Otoplasty

Otoplasty is done to correct deformity of the outer ear. Otoplasty may be done to bring the ears closer to the head and reduce the size of ears that are larger than normal. Unusually shaped ears may also be corrected. The doctor makes an incision behind the ear, and the normal cartilage folds and shape of the ears are recreated.

BEFORE SURGERY:

- Minimize aspirin, Motrin, Ibuprofen, Advil, Nuprin or any other anti-inflammatory medicine for one week before and one week after surgery. Tylenol is permitted at any time.
- If you take Coumadin or other blood thinners please discuss this with your surgeon.
- Please do not eat or drink anything after midnight the night before surgery, but you may take any
 prescription medicines the morning of surgery with a sip of water.
- The hospital or surgical center will call you the day before with time and instructions for surgery. Please make arrangements for a ride home after surgery, as you will not be able to drive.

PAIN MANAGEMENT AFTER SURGERY:

- There is usually moderate pain the first night after surgery, mostly form the bulky pressure dressing worn for the first 24 hours. After the bulky dressing is removed the next day, there typically isn't much pain after this surgery.
- The ears will be tender, so if the patient usually sleeps on their side, the sleeping pattern will be disrupted for a week or so following the surgery. Elevating the head of the bed or sleeping on a few pillows will help to alleviate some of the pain.
- Give pain medications such as Tylenol as necessary

WOUND CARE:

- A tight dressing will be placed over the ears to keep the ears from developing a blood clot and to decrease the swelling. The dressing will be removed one day after surgery and the ears will be examined.
- It is important to protect the ears against bending or folding for 6 weeks. A 'beanie' cap or an ear band like you would use in cold weather works the best. This head band should be worn day and night for 3 weeks, then at night only for 3 more weeks.

DRAINAGE:

- A small amount of blood/drainage is common, especially the first night.
- Soft drains are often placed behind the ears, and removed with the bulky dressing the day after surgery.
- The bloody drainage usually does not soak through the dressing completely, but more gauze can be applied over the dressing if necessary.
- The Physician must be the one to remove the dressing.
- Call your Doctor if the bleeding does not stop.

BATHING/SHOWERING:

• For the first week after surgery, the ear should not get wet, so be cautious with baths and showers. You may bathe as usual after one week.

ACTIVITY:

- Avoid sports or strenuous activities for 2-3 weeks.
- Contact sports and ball activities should be avoided for 6 weeks after surgery. There is a risk of deformity or loss of correction if a direct injury to the ear should occur.

WHEN TO CALL THE DOCTOR:

- Redness or tenderness of the ear
- Swelling
- Uncontrolled pain
- Temperature over 101.5 degrees Fahrenheit
- Excessive bleeding/drainage—please do not take aspirin or Motrin (Advil or Ibuprofen) for one week before or one week after surgery.
- Signs of dehydration—if the patient has had very little fluid intake for 12 hours or if a child seems particularly irritable or has not urinated for 8-10 hours.

RISKS:

- The ears will not be perfect, nor will they be perfectly symmetric, but they should be greatly improved.
- If a blood clot develops or the patient doesn't take the antibiotics, an infection could occur that could lead to cartilage loss, skin loss, unusual scarring, and a deformed ear.
- The shape of the ear canal could be deformed, resulting in cerumen impaction and hearing loss.
- One or both ears may slowly start pushing back out again, requiring more surgery. Small touch-up surgery can usually be done in the clinic with local anesthesia.

Please Call our office anytime with Questions or concerns, we are always happy to help!

Derrick Gale MD

Assistants:

Maty Fish - Medical Assistant

Phone: 801-328-2522 Extension 1409

Eliana Arellano - Patient Coordinator

Call or Text: 385-226-5577